
Reversing the Opioid Epidemic in Alabama


A HEALTH CARE PROFESSIONAL'S TOOLBOX TO REVERSE THE OPIOID EPIDEMIC



Despite the efforts of physicians, the public health community and elected officials, the opioid overdose and death epidemic continues in Alabama and across the nation. This toolbox contains important information and resources for physicians and health care professionals across the state.




Source: U.S. Centers for Disease Control and Prevention



PRESCRIPTIONS FOR:

Buprenophine, Buprenophine/Naloxone, Methadone, Naloxone, Naltrexone.







Reversing the Opioid Epidemic in Alabama

TREATING THE PATIENT'S PAIN WITH CARE AND COMPASSION

CHECK the state prescription drug monitoring program (PDMP) to ensure you have the information you need about your patient's prescription history.

AVOID initiating opioids for new patients with chronic non-cancer pain unless the expected benefits are anticipated to outweigh the risks. Non-pharmacologic therapy and non-opioid therapies are preferred.

LIMIT the amount of opioids prescribed for post-operative and acute care. It is recommended that patients only receive the lowest effective dose for the shortest possible duration.

DETERMINE the total daily amount of each opioid the patient takes.

CONVERT each to MMEsmultiply the dose for each opioid by the conversion factor (see table below)

ADD them together.

CALCULATING MORPHINE **MILLIGRAM EQUIVALENTS (MME)**

~	•		
OPIOID (Doses in mg/day except where noted)	CONVERSION FACTOR		
Codeine	0.15		
Fentanyl trans-dermal (mcg/hr)	2.4		
Hydrocodone	1		
Hydromorphone	4		
Methadone			
1-20 mg/day	4		
21-40 mg/day	8		
41-60 mg/day	10		
≥ 61-80 mg/day	12		
Morphine	1		
Oxycodone	1.5		
Oxymorphone	3		

These dose conversions are estimated and cannot account for all individual differences in genetics and pharmacokinetics.

MORPHINE MILLIGRAM **EQUIVALENT REFERENCE GUIDE**

DOSE PER DAY

DOSE PER DAY:	1	2	3	4	5
Tramadol	5	10	15	20	25
Codeine 30*	4.5	9	13.5	18	22.5
Codeine 60**	9	18	27	36	45
Hydrocodone 5	5	10	15	20	25
Hydrocodone 7.5	7.5	15	22.5	30	37.5
Hydrocodone 10	10	20	30	40	50
Oxycodone 5	7.5	15	22.5	30	37.5
Oxycodone 7.5	11.25	22.5	33.75	45	56.25
Oxycodone 10	15	30	45	60	75
Morphine 15	15	30	45	60	75
Morphine 30	30	60	90	120	150
Morphine 60	60	120	180	240	300
Dilaudid 4	16	32	48	64	80
Dilaudid 8	32	64	96	128	160
Fentanyl 12	30				
Fentanyl 25	60				
Fentanyl 50	120				
Fentanyl 100	240				
Nucynta 50	20	40	60		
Nucynta 75	30	60	90		
Nucynta 100	40	80	120		

Codeine 30* = Tylenol 3 Codeine 60** = Tylenol 4

GREY - check PDMP twice a year.

GREEN - check every prescription.

MME Calculations determined using OpioidCalc

CAUTION: Do not use the calculated dose in MMEs to determine dosage for converting one opioid to another—the new opioid should be lower to avoid unintentional overdose caused by incomplete cross-tolerance and individual differences in opioid pharmacokinetics. Consult the medication label.

USE EXTRA CAUTION: Methadone - the conversion factor increases at higher doses. Fentanyl - dosed in mcg/hr instead of mg/ day, and absorption is affected by heat and other factors.

WHAT DO ALABAMA BOARD OF **MEDICAL EXAMINERS REOUIRE?**

For controlled substance Rxs totaling 30 MME/day or less, physicians are expected to use the PDMP in a manner consistent with good clinical practice.

For controlled substance Rxs totaling more than 30 MME/day, physicians shall review the patient's Rx history in the PDMP at least two times per year.

For controlled substances more than 90 MME/day, physicians shall query the PDMP to review a patient's Rx history on the same day the Rx is written.

ALABAMA LAW ALLOWS:

A licensed physician approved by the ADPH who has authority to prescribe, dispense, or administer controlled substances may designate up to two employees who may access the PDMP on the physician's behalf.

RESOURCES:

Alabama Smart and Safe program: www.smartandsafeal.org

Alabama Prescription Drug Monitoring Program: www.adph.org/pdmp

Providers' Clinical Support System: https://pcssnow.org/educationtraining/treating-chronic-pain-core-curriculum/

CDC Guidelines for Prescribing Opioids for Chronic Pain: www.cdc.gov/ mmwr/volumes/65/rr/rr6501e1.htm

AL County-level map of opioid prescribing: CDC Morphine Milligram Equivalent information: https://www.cdc.gov/ drugoverdose/pdf/ calculating total daily dose-a.pdf







Reversing the Opioid Epidemic in Alabama

OVERDOSE PREVENTION

AMA TASK FORCE TO REDUCE OPIOID ABUSE RECOMMENDATION ON NALOXONE

CO-PRESCRIBING NALOXONE MAY HELP SAVE A PATIENT'S LIFE.

Several factors may be helpful in determining whether to co-prescribe naloxone to a patient, to a family member or to a close friend of the patient, including:

- Is my patient on a high dose of opioids?
- · Does my patient also have a concomitant benzodiazepine prescription?
- · Does my patient have a history of substance use disorder?
- · Does my patient have an underlying mental health condition that might make him or her more susceptible to overdose?
- Does my patient have a medical condition, such as a respiratory disease or other co-morbidities, that might make him or her susceptible to opioid toxicity, respiratory distress or overdose?
- Might my patient be in a position to aid someone who is at risk of opioid overdose?

LEARN MORE:

https://www.end-opioid-epidemic.org/ wp-content/uploads/2017/08/AMA-Opioid-Task-Force-naloxone-one-pager-updated-August-2017-FINAL-1.pdf

relat

Prescribe to Prevent's Naloxone Product Chart: http://prescribetoprevent.org/wp2015/wp-content/up loads/Naloxone-product-chart.17_04_14.pdf

WHAT DOES ALABAMA LAW AUTHORIZE?

Alabama law (Section 20-2-280)

authorizes physicians acting in good faith to directly or by standing order prescribe naloxone to:

An individual at risk of experiencing an opiate-related overdose.


A family member, friend, or other individual, including law enforcement, in a position to assist an individual at risk of experiencing an opiate-related overdose.

FDA requires strong warnings for opioid analgesics, prescription opioid cough products, and benzodiazepine labeling related to serious risks and death from combined use.

Learn more:

www.fda.gov/NewsEvents/Newsroom/Pres sAnnouncements/ucm518697.htm

PROMOTE SAFE STORAGE AND DISPOSAL OF OPIOIDS AND **ALL MEDICATIONS**



REMIND YOUR PATIENTS! Store medicines out of reach from children and never share prescriptions with anyone.

URGE YOUR PATIENTS TO SAFELY DISPOSE OF EX-PIRED, UNWANTED AND UNUSED MEDICATIONS! Recommend patients use pharmacy and law enforcement "take back" resources whenever possible.

Learn Where and How to Dispose of Unused Medicines:

https://www.end-opioid-epidemic.org/wp-content/uploads/2017/05/op ioid-safe-storage-and-disposal.pdf

Search for a disposal location near you:

www.fda.gov/downloads/ForConsumers/ConsumerUpdates/UCM449824.pdf





Reversing the Opioid Epidemic in Alabama

TREATMENT FOR OPIOID USE DISORDER

Stop the stigma of substance use disorder.


Enhance access to treatment.



"Patients in pain and patients with a substance use disorder need comprehensive treatment – not judgment. Just as a patient with heart disease or diabetes, a patient with a substance use disorder can be successfully treated and enjoy long-term recovery."

- Patrice A. Harris MD, MA, Chair, AMA Task Force to Reduce Opioid Abuse

TREATMENT, CONSULTATION, AND TRAINING RESOURCES



DOWNLOAD THE DIRECTORY: http://www.mh.alabama.gov/downloads/SA/SASDProgramDirectory.pdf



Become trained to provide in-office buprenorphine (PCSS) https://pcssnow.org/educationtraining/mat-training/

ASAM National Practice Treatment Guideline (PCSS-MAT) www.pcssmat.org/the-asam-na tional-practice-guideline-for-theuse-of-medications-in-the-treat ment-of-addiction-involving-opi oid-use

Providers' Clinical Support System for **Medication Assisted Treatment** www.pcssmat.org

Safe Medicine Disposal www.disposemymeds.org Opioid Treatment Program Directory dpt2.samhsa.gov/treatment/ direc tory.aspx

Physician Locator for Substance Use Disorder Treatment (SAMHSA) www.samhsa.gov/medication-as sisted-treatment/physician-pro gram-data/treatment-physi cian-locator? eld_bup_ physician/ us_state_value=AL

SAMHSA MAT Free Pocket Guide http://www.store.samhsa.gov/product/ Medication-Assist ed-Treatment-of-Opioid-Use-Dis order-Pocket- Guide/Most-Popular/ SMA16- 4892PG?sortByValue=4

Treatment Near You www.zeroaddiction.org/nd-help



Funding for this initiative was made possible (in part) by grant no. 5H79TI025595 from SAMHSA. The views expressed in written conference materials or publications and by speakers and moderators do not necessarily reflect the official policies of the Department of Health and Human Services; nor does mention of trade names, commercial practices, or organizations imply endorsement by the U.S. Government.



